

Drug Facts

Active ingredient (in each caplet)

Calcium Polycarbophil 625 mg

(equivalent to 500 mg polycarbophil).....Bulk-forming laxative

Purpose

- Uses** • relieves occasional constipation • promotes regularity
• this product generally produces a bowel movement in 12 to 72 hours

Warnings

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. • Do not take this product if you have difficulty in swallowing. • If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Do not use • for more than 1 week unless directed by a doctor

Ask a doctor before use if you have • abdominal pain, nausea, or vomiting
• a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are • taking any other drug
• all laxatives may affect how other drugs work.

Stop use and ask a doctor if • you have rectal bleeding • you have no bowel movement within 3 days • these could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

GERI CARE

NDC 57896-449-09

Fiber Laxative

Calcium Polycarbophil

- Relieves Constipation
- Promotes Regularity

Compare to active ingredient
in FiberCon®*

90 Caplets • 625 mg Each

Directions • do not exceed recommended dose • take each dose with at least 8 ounces (a full glass) of fluid; taking this product without enough liquid may cause choking; see choking warning • do not take this product within 2 hours before or after taking any other drug

adults and children 12
years of age and older

Starting dose: 2 caplets once a day
Maximum dose: 2 caplets four times a day
or as directed by a doctor

children under 12 years of age

consult a doctor

Other information • each caplet contains: calcium 138 mg

• store at controlled room temperature, 68°F-77°F (20°C-25°C)

• protect contents from moisture • for institutional use only

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Inactive ingredients: caramel, cellulose, crospovidone, hypromellose, magnesium silicate, magnesium stearate, mineral oil, PEG, silica, sodium lauryl sulfate, stearic acid. May also contain calcium carbonate, croscarmellose sodium, maltodextrin, talc.

*This product is not manufactured or distributed by the owner of the registered trademark FIBERCON®.
REV 0515RZC

Dist. by: Geri-Care Pharmaceuticals Corp.
1650 63rd Street, Brooklyn, NY 11204

