GERICARE

NDC 57896-921-01

Regular Strength Enteric Coated Aspirin

Pain Reliever (NSAID)

PACKAGE NOT CHILD RESISTANT

100 Tablets 325 mg each

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 325 mg (NSAID)*	Pain reliever

*nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains ■ ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have: ▫ high blood pressure ▫ heart disease ▫ liver cirrhosis ▫ kidney disease ▫ asthma

Distributed By: Geri-Care Pharmaceuticals Corp.
1650 63rd Street, Brooklyn, NY 11204

REV 0113ALT

3 57896 92101 9

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are ■ taking any other drug containing an NSAID (prescription or nonprescription) ■ taking a blood thinning (anticoagulant) or steroid drug ■ taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ▫ feel faint ▫ vomit blood ▫ have bloody or black stools ▫ have stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away. ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur ■ ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- swallow whole, do not chew or crush
- do not exceed recommended dose

adults and children 12 years and older	take 1-2 tablets every 4 hours, as needed, not more than 12 tablets in 24 hours, or as directed by a doctor
children under 12	ask a doctor

Other information ■ store at room temperature 15°C - 30°C (59°F - 86°F) ■ for institutional use only

Inactive ingredients: cellulose, D&C yellow #10 lake, FD&C yellow #6 lake, hypromellose, iron oxide, PEG, polydextrose, polyvinyl acetate phthalate, propylene glycol, shellac wax, silica, simethicone, sodium alginate, sodium bicarbonate, starch, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate. May also contain acetylated monoglycerides, croscarmellose sodium, hypromellose phthalate, methacrylic acid, mineral oil, polysorbate 80, sodium lauryl sulfate.

STOP PEELING