## **Usage Instructions**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all directions and precautions on this product label. Not recommended for use on unfinished, oiled, or waxed wood surfaces. For other surfaces, test in an inconspicuous area. TO DISINFECT: Hard, non-porous non-food contact surfaces: Pre-clean Surface. Hold container 6"-8" from surface and spray until thoroughly wet. Bacteria: Let stand for 60 seconds. Wipe with a cloth. Bacteria, Viruses and Fungi: Let stand for 5minutes. Wipe with a cloth. SANITIZING: Hold container 6"-8" from surface and spray until thoroughly wet. To Sanitize Hard Non-porous non-food contact surfaces: Let stand 10 seconds. Wipe clean with a damp cloth. Pre-clean heavily soiled surfaces. To Sanitize Hard Non-porous non-food contact surfaces FOR 24 HOURS: - Let stand 5 minutes. Allow to air dry. Pre-clean heavily soiled surfaces: Let stand for 60 seconds. Let air dry. Pre-clean heavily soiled surfaces. To DEODORIZE: Spray on surfaces as needed. TO CLEAN: Spray on surfaces, allow to penetrate. Wipe clean with a cloth. TO PREVENT MOLD AND MILDEW growth: On Fabrics: Apply to fabric surface until wet. Allow to air dry. Repeat every 28 days to inhibit mold and mildew growth. Pre-clean heavily soiled surfaces. On hard surfaces: Thoroughly wet surface. Allow to air dry. Repeat every 7 days to inhibit mold and mildew growth. Pre-clean heavily soiled surfaces. STORAGE AND DISPOSAL: Store in original container in areas inaccessible to small children. Do not puncture or incinerate. Place in trash or offer for recycling.

## Suggested Use

KILLS IN 5 MINUTES: Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus, Escherichia coli O157:H7, Enterobacter aerogenes, Listeria monocytogenes, Methicillin Resistant Staphylococcus aureus - MRSA, Streptococcus pyogenes, †Human Coronavirus, †Herpes Simplex virus type 1, †Herpes Simplex virus type 2, †Influenza A H1N1, †Respiratory Syncytial Virus [RSV], †Rotavirus, †Rhinovirus, †Norovirus (Feline Calicivirus as surrogate), Trichophyton mentagrophytes, Aspergillus niger