

2657 Anbesol Maximum Strength Oral Anesthetic

Active ingredient

Benzocaine 20% - Oral anesthetic

Inactive ingredients

benzyl alcohol, carbomer 934P, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, glycerin, methylparaben, natural and artificial flavor, polyethylene glycol, propylene glycol, saccharin

Uses

Temporarily relieves pain associated with the following mouth and gum irritations:

- toothache
- sore gums
- canker sores
- braces
- minor dental procedures
- dentures

Warnings

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics

When using this product:

- avoid contact with the eyes
- do not exceed recommended dosage
- do not use for more than 7 days unless directed by a doctor/dentist

Stop use and ask a doctor if:

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling, rash, or fever develops

Keep out of reach of children.

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions

To open tube, cut tip of the tube on score mark with scissors.

- adults and children 2 years of age and older: apply to the affected area up to 4 times daily or as directed by a doctor/dentist
- children under 12 years of age: adult supervision should be given in the use of this product
- children under 2 years of age: consult a doctor/dentist

For denture irritation:

- apply thin layer to the affected area
- do not reinsert dental work until irritation/pain is relieved
- rinse mouth well before reinserting

Other information:

Store at 20-25°C (68-77°F). Do not refrigerate.