



NDC 57896-226-01  
THE SENSIBLE CHOICE  
FOR PRICE AND HEALTH

# Extra Strength Pain-Reliever

Acetaminophen 250 mg  
Aspirin 250 mg (NSAID)  
Caffeine 65 mg

COMPARE TO THE ACTIVE  
INGREDIENTS IN EXCEDRIN®\*

## 100 Caplets

**Tamper Evident:** Do not use if imprinted seal under cap is missing or broken.

### Drug Facts

#### Active ingredients (in each caplet)

Acetaminophen 250 mg ..... Pain reliever  
Aspirin 250 mg (NSAID) ..... Pain reliever  
Caffeine 65 mg .....  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ Temporarily relieves minor aches and pains

#### Warnings

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than directed ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

\*This product is not manufactured or distributed by the owner of the registered trademark Excedrin®.

Distributed By: Geni-Care Pharmaceuticals Corp.  
1650 63rd Street, Brooklyn, NY 11204



3 57896 22601 5

PEEL HERE FOR MORE DRUG FACTS

### Drug Facts (continued)

**Do not use** ■ If you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use** ■ If you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use if you are taking** ■ any other drug containing an NSAID (prescription or nonprescription) ■ a blood thinning (anticoagulant) or steroid drug ■ a prescription drug for diabetes, gout, or arthritis ■ any other drug, or are under a doctor's care for any serious condition

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away.

**You experience any of the following signs of stomach bleeding:** ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely decided to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

**Overdose Warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not use more than directed (see overdose warning)

■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours  
■ children: under 12: ask a doctor

#### Other information

■ store at room temperature 15° - 25°C (59° - 77°F)

**Inactive ingredients** carmelumina wax, cellulose, hypromellose, mineral oil, povidone, silica, sodium starch glycolate, starch, stearic acid, titanium dioxide.

STOP PEELING