

4015 Non-Aspirin Extra-Strength Pain Relief

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg Pain Reliever/Fever Reducer

Purposes

Uses

- Temporarily relieves minor aches and pains
 - Temporarily reduces fever
-

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- More than 8 caplets in 24 hours, which is the maximum daily amount
 - With other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product
-

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if:

- Pain gets worse or lasts more than 10 days
- Fever gets worse or lasts more than 3 days
- New symptoms occur
- Redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children

Overdose Warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:

- **Do not take more than directed**
 - Adults and children 12 years and over: take 1-2 caplets every 4-6 hours as needed; not more than 8 caplets in 24 hours
 - Children under 12 years: do not use
-

Other Information

- **TAMPER EVIDENT:** Do not use if imprinted seal under cap is missing or broken
 - Store at 20°C - 25°C (68°F - 77°F)
 - For institutional use only
-

Inactive Ingredients:

Povidone, sodium starch glycolate, starch, stearic acid. May also contain: crospovidone, methylparaben and propylparaben
