4015 Non-Aspirin Extra-Strength Pain Relief

Drug Facts

Uses

- Temporarily relieves minor aches and pains
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Sever liver damage may occur if you take:

- More than 8 caplets in 24 hours, which is the maximum daily amount
- With other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if:

- Pain gets worse or lasts more than 10 days
- Fever gets worse or lasts more than 3 days
- New symptoms occur
- Redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children

Overdose Warning: Taking more than the recommended does (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:

- Do not take more than directed
- Adults and children 12 years and over: take 1-2 caplets every 4-6 hours as needed; not more than 8 caplets in 24 hours
- Children under 12 years: do not use

Other Information

- TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken
- Store at 20°C 25°C (68°F 77°F)
- For institutional use only

Inactive Ingredients:

Povidone, sodium starch glycolate, starch, stearic acid. May also contain: crospovidone, methylparaben and propylparaben