

4139 Milk of Magnesia 16oz

Drug Facts

Active ingredient (in each 15mL tablespoonful)	Purpose
Magnesium hydroxide 1200 mg	Saline laxative

Use

- Relieves occasional constipation (irregularity)
 - Usually produces bowel movement in ½ to 6 hours
-

Warnings

Ask a doctor before use if you have:

- Kidney disease
 - A magnesium-restricted diet
 - Stomach pain, nausea, or vomiting
 - A sudden change in bowel habits that lasts over 14 days
-

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if:

- You have rectal bleeding or no bowel movement after using this product.
These could be signs of a serious condition
 - You need to use a laxative for more than one week
-

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- Shake well before use
 - Do not exceed the maximum recommended daily dose in a 24 hour period
 - Dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
 - Drink a full glass (8oz) of liquid with each dose
-

Adults and children 12 years and older	2 to 4 tablespoonfuls (TBSP)
Children 6 to 11 years	1 to 2 tablespoonfuls (TBSP)
Children under 6 years	Ask a doctor

Other information

- Each 15 mL tablespoonful contains: magnesium 500 mg
 - Store at room temperature and avoid freezing
 - Close cap tightly after use
 - **TAMPER-EVIDENT:** Do not use if imprinted neckband is missing or broken.
-

Inactive Ingredients

purified water, sodium hypochlorite
