4172 Stool Softener 100 mg

Drug Facts

Active ingredient (in each softgel)

Purpose

Docusate Sodium 100 mg..... Stool Softener Laxative
Uses

- Relieves occasional constipation (irregularity)
- Generally produces bowel movement in 12 to 72 hours

Warnings

Ask a doctor before use if you

- Have stomach pain, nausea or vomiting
- Have sudden change in bowel habits that persists over a period of 2 weeks
- Are presently taking mineral oil

Stop use and ask a doctor if

- You need to use a laxative longer than one week
- You have rectal bleeding or fail to have a bowel movement

These could be signs of a serious condition

If pregnant or breastfeeding, ask a health professional before use.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- Adults and children 12 years an dolder : take 1-2 softgels daily until first bowel movement; 1 softgel daily thereafter or as directed by doctor
- Children under 12: consult a doctor
- Do not exceed recommended dose

Other Information

- Each softgel contains: sodium 5 mg. Very low sodium.
- Store at 59°-77°F (15°-25°C)
- Keep tightly closed
- Product from USA or Canada
- Tamper Evident: Do not use if imprinted seal under cap is missing or broken

Inactive Ingredients: FD&C red #40, gelatin, glycerin, edible ink, PEG, propylene glycol, sorbitol special, water. May also contain D&C yellow #10, FD&C yellow #6 (sunset yellow), mannitol.