## 

runny nosesneezing

itchy, watery eyes
 itching of the nose or throat

## Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast feeding, ask a health professional before

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

0 T 11 ( 10 T 1

THE STATE OF THE S

NDC 57896-788-09
THE SENSIBLE CHOICE
FOR PRICE AND HEALTH

## 24 Hour Non-Drowsy\* Allergy Relief **Loratacline**

Antihistamine

Compare to active ingredient in Claritin®\*

When taken as directed. See Drug Facts panel

Directions

adults and children 12 years and over than 1 tablet daily; not more than 1 tablet in 24 hours children under 12 years of age ask a doctor consumers with liver or kildney disease! ask a doctor

Other information • TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken. • store between 2°C and 30°C (36F-86F) • product of Canada • protect from excessive moisture Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose Puestions? call 1-800-540-3765

\*This product is not manufactured or distributed by the owner of the registered trademark Claritin®. REV 0413A

DIST. BY: GERI-CARE PHARMACEUTICALS CORP., 1650 63rd St., Brooklyn, New York 11204

3 57896 78809

90 Tablets 10 mg Each