5970 Female Catheter Kit Instructions and Ingredients

Directions:

Contents arranged in sequence:

- 1. Unfold CSR wrap and use as a sterile field
- 2. Don gloves
- 3. Pull catheter out of tube to desired length. Lay tube on sterile field.
- 4. Prepare patient and proceed with catheterization. (**Note**: If urine does not flow freely into the specimen tube, loosen the cap slightly.)
- 5. After sample collection, pull catheter out of top; tighten cover and depress spout.
- 6. Send to lab in normal manner.

Aplicare Three Povidone-Iodine Swabsticks

| Drug Facts | |
|-------------------------|------------|
| Active ingredient | Purpose |
| Povidone-iodine USP 10% | Antiseptic |

Use

Antiseptic skin preparation

Warnings

Do not use:

- If allergic to iodine
- In the eyes

For external use only

Ask a doctor before use if injuries are:

- Deep or puncture wounds
- Serious burns

Stop use and ask a doctor if:

- Redness, irritation, swelling, or pain persists or increases
- Infection occurs

Avoid pooling beneath patient

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or consult a poison control center immediately.

Directions

Apply locally as needed

Other information

- 1% titratable iodine
- Latex free
- For hospital or professional use only
- STERILE unless opened or damaged

Inactive ingredients

Citric acid, disodium phosphate, nonoxynol-9, sodium hydroxide, water

Dover Lubricating Jelly 5g

- Water-soluble lubricant.
- For hospital and professional use.
- Will not harm surgical instruments, rubber, or plastic.
- Non-irritating to skin, tissue, and mucous membranes.
- Do not use if package is opened or damaged.